

## **Instructions for Medications:**

You have been prescribed two types of pain medications

1<sup>st</sup> pain Medication is a Narcotic  
(Tylenol #3, Vicoden Es, Percocet)

These medications cause drowsiness, nausea (on an empty stomach)

**-You can not take this medication and operate a motor vehicle.**

Directions:

Take the narcotic medicine every 4 to 6 hours as needed for pain. **WITH FOOD!!!**

2<sup>nd</sup> pain Medication is Motrin 800mgs (Ibuprofen) it is equal to four Advil's.  
This type of medication will not cause drowsiness; you can take this and operate a motor vehicle, work, etc.

Directions:

1 tablet every 4 to 6 hours between your doses of the Narcotic.

\*If you experience severe pain and the narcotic is not helping, you can take the Motrin and the Narcotic at the same time. **DO NOT** take two narcotics at the same time. This can lead to respiratory depression and possible death.

Your Doctor sometimes prescribes an anti-nausea medication called Compazine. This medicine does cause drowsiness, and should only be taken if the patient is nauseous and or vomiting.

An antibiotic may be prescribed to you the antibiotic should be taken the day **AFTER** the procedure, taking one in the AM and one in the PM. Ideally with breakfast and one with dinner, usually lasting five days unless otherwise prescribed.

The Doctor as well as the staff will review this with you. If you have any questions please do not hesitate to ask.